

REPTILIA: SQUAMATA: SERPENTES: COLUBRIDAE

ALSOPHIS ANOMALUS

Catalogue of American Amphibians and Reptiles.

Powell, R. and R.W. Henderson. 1998. *Alsophis anomalus*.

***Alsophis anomalus* (Peters)**
Hispaniolan Racer

Zamenis anomalus Peters 1863:282. Type locality, unknown. Holotype, Museum für Naturkunde, Humboldt Universität, Berlin (ZMB) 2269; sex, collector, and date of collection unknown (not examined by authors).

Dromicus anomalus: Fischer 1888:37.

Alsophis anomalus: Barbour 1914:336. First use of present combination.

Leimadophis anomalus: Amaral 1929 (1930):164.

• **Content.** No subspecies are recognized.

• **Definition.** This large (maximum SVL to 1770 mm) colubrid has 197–215 ventrals, 109–130 subcaudals, 21 dorsal scale rows at midbody, 7–9 (mode 8) supralabials, 10–12 infralabials, 1/1 preoculars, 2/2 postoculars, 1/1 large suboculars, 1+2/1+2 temporals, and 1/1 elongate loreals. Dorsal scales are smooth and the cloacal scute (= anal plate) is divided.

The dorsum is brown, often with a bronze cast, and usually is patternless in adults, although specimens from Île de la Tortue may exhibit a faint pattern of anterior chevrons replaced posteriorly by crossbars. In adults, the free edges of the dorsal scales are often darker than the rest of the scale. The venter may be yellow or beige, without markings or with scattered, irregular dark blotches. The mental, chin shields, gulars, and infralabials are gray, and this gray wash may extend posteriorly on the free edges of anterior ventrals. The juvenile dorsal pattern consists of ragged crossbands 1–3 scales wide (only 1 scale wide on tail); the venter (body and tail) is immaculate ivory with a yellowish wash on some portions.

• **Diagnosis.** *Alsophis anomalus* can be distinguished from other Hispaniolan colubrids by its 21 dorsal scale rows at midbody. The only sympatric congener, *A. melanichnus*, has 17 dorsal scale rows at midbody. All other Hispaniolan colubrids either have extremely slender bodies and 13 or fewer posterior dorsal scale rows (*Uromacer*) or have 19 dorsal scale rows at midbody (*Antillophis*, *Darlingtonia*, *Hypsirhynchus*, *Ialtris*).


• **Descriptions.** Detailed descriptions are in Cochran (1941) and Schwartz and Henderson (1991).

• **Illustrations.** Line drawings of the head are in Cochran (1941).

• **Distribution.** A Hispaniolan endemic, this species has a wide but disjunct distribution in dry areas throughout the western half of the island. Specimens also have been taken on Île de la Tortue and Isla Beata. The range has been illustrated in Schwartz and Henderson (1991).

• **Fossil Record.** None.

• **Pertinent Literature.** Maglio (1970) discussed origin, phylogeny, and zoogeography. Schwartz (1980) characterized *A. anomalus* as an islandwide form. Ducorday (1981) discussed the ecology of *A. anomalus* on Isla Beata. Henderson and Sajdak (1986) noted distribution in relation to that of the mongoose. SEA/DVS (1990) provided an index to habitats in the Dominican Republic. Henderson and Sajdak (1996) provided informa-



Map. Range of *Alsophis anomalus* (modified from Schwartz and Henderson 1991). The type locality is too imprecise to plot, dots mark known records.



Figure 1. *Alsophis anomalus* from Palmiste, Île de la Tortue, Haiti. Photograph courtesy of C. Rhea Warren.




Figure 2. Head of an adult female *Alsophis anomalus* (USNM 59917) from Jean Rabel, Haiti (from Cochran 1941). The left subocular on this snake was fused with the fifth supralabial.

tion on diet. Rodríguez-Robles and Greene (1996) included this species in their study of ecological relationships of Greater Antillean snakes.

The species is included in notes, checklists, guides, and keys by Fischer (1888), Müller (1892), Boulenger (1894), Meerwarth (1901), Amaral (1929 [1930]), Barbour (1930, 1935, 1937), Cochran (1924, 1934), Schwartz and Thomas (1975), Henderson and Schwartz (1984), Henderson et al. (1984), Schwartz and Henderson (1985, 1988), Franz and Cordier (1986), Henderson and Crother (1989), and Powell et al. (1996).

Frank and Ramus (1995) proposed the common name "Hispaniola [*sic*] racer" for this species.

• **Etymology.** The name *anomalus* is derived from the Greek *anomalos* (uneven or irregular), possibly in reference to the faint dorsal pattern elements in these snakes.

• **Comment.** The rarity of this species is inexplicable; a fresh roadkill found in the Valle de Neiba in June 1997 (Bobby Witcher Memorial Collection, Avila College, Kansas City, MO 64145 [BWMC] 06124) was the first main island specimen taken in many years. Apparently suitable habitats are widespread and populations of other Hispaniolan colubrids often are very dense compared with those of most Neotropical snakes.

Literature Cited

- Amaral, A. 1929 (1930). Estudos sobre ophidios neotropicos. XVIII. Lista remissiva dos ophidios da região neotropical. Mem. Inst. Butantan 4:129–271.
- Barbour, T. 1914. A contribution to the zoogeography of the West Indies, with especial reference to amphibians and reptiles. Mem. Mus. Comp. Zool. 44:205–359.
- . 1930. A list of Antillean reptiles and amphibians. Zoologica (N.Y.) 11:61–116.
- . 1935. A second list of Antillean reptiles and amphibians. Zoologica (N.Y.) 19:77–141.
- . 1937. Third list of Antillean reptiles and amphibians. Bull. Mus. Comp. Zool. 82:77–166.
- Boulenger, G.A. 1894. Catalogue of the Snakes in the British Museum (Natural History). Vol. 2. Trustees of the British Museum, London.
- Cochran, D.M. 1924. Notes on the herpetological collections made by Dr. W.L. Abbot on the island of Haiti. Proc. U.S. Natl. Mus. 66:1–15.
- . 1934. Herpetological collections made in Hispaniola by the Utowana Expedition, 1934. Occ. Pap. Boston Soc. Nat. Hist. 8:163–188.
- . 1941. The herpetology of Hispaniola. Bull. U.S. Natl. Mus. (177):vii + 398 p.
- Ducoudray, F. S. 1981. La isla de la serpiente, p. 137–145. In Investigaciones en las Islas Beata y Alto Velo. Ser. Invest. Antropol. 18. Mus. Hombre Dominicano y Marina de Guerra, Santo Domingo.
- Fischer, J.G. 1888. Über eine Kollektion Reptilien und Amphibien von Hayti. Jahrb. Wiss. Anst. Hamburg 5:23–45.
- Frank, N. and E. Ramus. 1995. A Complete Guide to Scientific and Common Names of Reptiles and Amphibians of the World. NG Publ., Inc., Pottsville, Pennsylvania.
- Franz, R. and D. Cordier. 1986. The Herpetofauna of the Proposed National Parks in Southern Haiti. USAID/Haiti Rept., Univ. Florida, Gainesville.
- Henderson, R.W. and B.I. Crother. 1989. Biogeographic patterns of predation in West Indian colubrid snakes, p. 479–517. In C.A. Woods (ed.), Biogeography of the West Indies: Past, Present, and Future. Sandhill Crane Press, Gainesville, Florida.
- and R.A. Sajdak. 1986. West Indian racers: a disappearing act or a second chance? Lore 36(3):13–18.
- and —. 1996. Diets of West Indian racers (Colubridae: *Allophis*): composition and biogeographic implications, p. 327–338. In R. Powell and R.W. Henderson (eds.), Contributions to West Indian Herpetology: A Tribute to Albert Schwartz. SSAR Contrib. Herpetol. Vol. 12. Ithaca, New York.
- and A. Schwartz. 1984. A guide to the identification of the amphibians and reptiles of Hispaniola. Milwaukee Pub. Mus. Spec. Publ. Biol. Geol. (4):1–70.
- , —, and S.J. Incháustegui. 1984. Guía para la indentificación de los anfibios y reptiles de la Hispaniola. Mus. Nac. Hist. Nat. Ser. Mono. (1):1–128.
- Maglio, V.J. 1970. West Indian xenodontine colubrid snakes: their probable origin, phylogeny, and zoogeography. Bull. Mus. Comp. Zool. 141:1–53.
- Meerwarth, H. 1901. Die westindischen Reptilien und Batrachier des naturhistorischen Museums in Hamburg. Mitt. Naturhist. Mus. Hamburg 18:1–41.
- Müller, F. 1892. Siebenter Nachtrag zum Katalog der herpetologischen Sammlung des Basler Museums. Festschr. 75, Verh. Naturf. Ges. Basel 10, pt. 1:195–215.
- Peters, W. 1863. Über einige neue oder weniger bekannte Schlangenarten des zoologischen Museums zu Berlin. Monatsb. Königl. Akad. Wiss. Berlin 1863:272–289.
- Powell, R., R.W. Henderson, K. Adler, and H.A. Dundee. 1996. An annotated checklist of West Indian amphibians and reptiles, p. 51–93. In R. Powell and R.W. Henderson (eds.), Contributions to West Indian Herpetology: A Tribute to Albert Schwartz. SSAR Contrib. Herpetol. Vol. 12. Ithaca, New York.
- Rodríguez-Robles, J.A. and H.W. Greene. 1996. Ecological patterns in Greater Antillean macrostomatan snake assemblages, with comments on body-size evolution in *Epicrates* (Boidae), p. 339–357. In R. Powell and R.W. Henderson (eds.), Contributions to West Indian Herpetology: A Tribute to Albert Schwartz. SSAR Contrib. Herpetol. Vol. 12. Ithaca, New York.
- Schwartz, A. 1980. The herpetogeography of Hispaniola, West Indies. Stud. Fauna Curaçao Carib. Isl. 61:86–127.
- and R.W. Henderson. 1985. A Guide to the Identification of the Amphibians and Reptiles of the West Indies Exclusive of Hispaniola. Milwaukee Pub. Mus., Milwaukee, Wisconsin.
- and —. 1988. West Indian amphibians and reptiles: a checklist. Milwaukee Pub. Mus. Contr. Biol. Geol. (74):1–264.
- and —. 1991. Amphibians and Reptiles of the West Indies: Descriptions, Distributions, and Natural History. Univ. Florida Press, Gainesville.
- and R. Thomas. 1975. A check-list of West Indian amphibians and reptiles. Carnegie Mus. Nat. Hist. Spec. Publ. (1): 1–216.
- SEA/DVS (Secretaría de Estado de Agricultura/Departamento de Vida Silvestre). 1990. La Diversidad Biológica en la República Dominicana: Reporte Preparado por el Departamento de Vida Silvestre para el Servicio Alemán de Cooperación Social-Técnica y Fondo Mundial para la Naturaleza (WWF-US). Apéndice. Sec. Estado Agric., SURENA/DVS. Sto. Domingo, República Dominicana.

Robert Powell, Department of Natural Sciences, Avila College, Kansas City, MO 64145, and **Robert W. Henderson**, Section of Vertebrate Zoology, Milwaukee Public Museum, Milwaukee, WI 53233.

Primary editor for this account, Larry David Wilson.

Published 30 January 1998 and Copyright © 1998 by the Society for the Study of Amphibians and Reptiles.
